


The Translator.



Computes all.

All there is to compute. Directly onto tape. Blue Shield enrollment information. Medicare information. Information information. The Translator means time saved in transcription. Time saved in key punch. Time saved for you. So in 1970, you'll have more time to be a doctor.



GREATER NEW YORK'S

BLUE SHIELD

UNITED MEDICAL SERVICE, INC.

Service is what we're all about.

The Bulletin of the New York Academy of Medicine, Vol. 47, No. 1, January 1971. Published monthly by the New York Academy of Medicine, 2 East 103 Street, New York, N. Y. 10029. Entered as second-class matter February 3, 1928, at the Post Office at New York, N. Y., under the act of August 24, 1912. Postage paid at New York, N. Y. Annual subscription United States \$15.00. Canada \$16.00. All other countries \$17.00. Single copies \$3.00.

Author and Subject Catalogs

LIBRARY OF THE NEW YORK ACADEMY OF MEDICINE

At the end of 1967 fully catalogued items in the Library of the New York Academy of Medicine included 365,000 bound volumes and 165,000 pamphlets, as well as portraits and other nonbook materials. Included also are the cards for thousands of foreign medical theses and several thousand bound volumes which have been sent to the Medical Library Center of New York.

In addition to the entries for journals, monographs, government documents, historical materials, and reference tools, the catalogs contain an extraordinary amount of additional information in analytics. Here may be found the contents of the important German *Handbücher* and *Festschriften*, journal supplements, and issues devoted to a single subject, proceedings of congresses and symposia of substance which might be published also as separate publications, and small notices whose information would not otherwise be available.

In the subject catalog may be found the Edwin Smith Surgical Papyrus, about 4,000 years old, and all library materials issued through December 31, 1967. The author-title catalog includes 1968 publications as well.

Author Catalog

Estimated 770,000 cards reproduced in 43 volumes

Price: \$2950.00

Subject Catalog

Estimated 610,000 cards reproduced in 34 volumes

Price: \$2400.00

10% additional charge on orders outside the U.S.

*Descriptive material on these titles and a complete catalog of publications
are available on request.*

G. K. HALL & CO. 70 Lincoln Street, Boston, Massachusetts 02111

Your Patients Deserve The Best



metropolitan pathology laboratory inc.

DIRECTED BY PHYSICIANS

MetPath is directed with one basic testing philosophy: inaccurate tests are worse than no tests at all. Laboratory Director is Dr. Joseph E. O'Brien, a graduate of the New York State University School of Medicine, and Board Certified in Anatomic and Clinical Pathology. Prominent consultants in academic medicine work actively along with our division directors, all of whom hold doctorate degrees, to assure that all diagnostic services performed continue to meet the physician's demand for quality laboratory service.

RIGID QUALITY CONTROL

Metropolitan Pathology Laboratory's rigid quality control uses automated procedures and computer-generated reports under the supervision of highly trained technicians to assure an extremely high degree of accuracy. Your inspection of our facilities is invited at any time.

PROFICIENCY TESTING

The excellence of a laboratory's work should be constantly checked. MetPath is engaged in a continuing proficiency program, using U.S. Public Health Service criteria. A copy of our latest proficiency rating will be sent upon request. As with all superior laboratories, there is no charge for repeat specimens when you, or our laboratory, feel the results do not coincide with the clinical diagnosis.

FEDERALLY LICENSED, MEDICARE APPROVED

MetPath is one of the less than ten percent of the nation's laboratories that are licensed under the rigid standards of the Clinical Laboratory Improvement Act, effective January 1st, 1969. MetPath has been approved for Medicare patients.

ONE-DAY TURNAROUND SERVICE IN PROTECTIVE CONTAINERS

Virtually all specimens received in the morning are processed the same day. Abnormal results are immediately telephoned to your office. Complete written reports follow within 24 hours. An inventory of mailing containers and specimen collection supplies is sent to you without charge.

LOWER PRICES

The prices we charge reflect the built-in economies that result from the use of advanced and highly automated equipment and large-scale testing procedures. We believe the prices of our services are lower than the prices of any laboratory of comparable quality.



metropolitan pathology laboratory inc.

185 West Englewood Avenue
Teaneck, New Jersey 07666
N.J. (201) 837-6440
N.J. (201) 488-1100
N.Y. (212) 928-5253

ROUTINE — ESOTERIC — HEALTH PROFILE SERVICES

ANNUAL HEALTH PROFILE

CHEMISTRY

ALBUMIN
ALKALINE PHOSPHATASE
BILIRUBIN
CALCIUM
CHOLESTEROL
GLUCOSE
LACTIC DEHYDROGENASE
PHOSPHORUS
SGO-TRANSAMINASE

TOTAL PROTEIN
UREA NITROGEN
URIC ACID
PROTEIN BOUND IODINE

MICROBIOLOGY

SEROLOGICAL SCREEN (V.D.R.L. or A.R.T.)
F.T.A. (performed on positive serological screen)

HEMATOLOGY

DIFFERENTIAL COUNT*
HEMATOCRIT**
HEMOGLOBIN
PERIPHERAL SMEAR***
RED BLOOD CELL COUNT
RED CELL INDICES**
WHITE BLOOD CELL COUNT

OBSTETRICS HEALTH PROFILE

HEMATOLOGY

ANTIBODY SCREEN (all patients regardless of Rh type)
ANTIBODY TITER (all patients with positive antibody screen)
BLOOD GROUP AND Rh TYPE
DIFFERENTIAL COUNT*
HEMATOCRIT**
HEMOGLOBIN
PERIPHERAL SMEAR***
RED BLOOD CELL COUNT
RED CELL INDICES**
WHITE BLOOD CELL COUNT

CHEMISTRY

BLOOD SUGAR (preferably 2 hour p.p.)
BLOOD UREA NITROGEN

CYTOLOGY

PAP SMEAR

MICROBIOLOGY

PREGNANCY TEST (when requested)
RUBELLA H. I.
SEROLOGICAL SCREEN (V.D.R.L. or A.R.T.)
F.T.A. (performed on positive serological screen)

*A complete differential count will be performed, at no additional charge, if any abnormality is noted in the hemoglobin, white blood count or peripheral smear.

**To maintain quality results this test is only performed on specimens received within 24 hours.

***If the specimen is to be referred by mail, the smear must be made at the time the blood is collected because blood cell morphology is altered by continuous contact with the anticoagulant.

The negative power of anxiety...

**This man thinks he may
never work again.**



The patient who has had a myocardial infarction is usually advised by his physician to avoid emotional excitement. All too often his family, acutely concerned, transmits its anxiety to him, urging him to "rest, rest."

How anxiety may interfere: In a study of 336 males who had suffered at least one myocardial infarction, Sigler¹ reports that manual workers showed the lowest percentage of patients returning to work, compared to clerical workers, business and professional men. The author notes that in many cases the mere apprehension that "return to work would shorten life prevents the patient from resuming activities." It is also well known that emotional disturbance is probably the most common cause of cardiac disability in postinfarction cases.¹

The anxiety factor in both *coronary* and *precoronary* patients has recently been discussed by Thomas,² who suggests: "Intensive investigation of the sources and kinds of anxiety, and how destructive forms of anxiety can be identified and relieved may be the next important step in the prevention of coronary heart disease."

Relief of anxiety with Librium® (chlordiazepoxide HCl) often proves a valuable adjunct to medical counsel, reassurance and the total management program; may help prevent the postcoronary patient from regressing into a state of invalidism.

As an adjunct in cardiovascular therapy, Librium® (chlordiazepoxide HCl): Quickly relieves anxiety of mild to severe degree in most cases. Helps expedite cooperation in therapeutic regimen. May be used concomitantly with certain specific medications of other classes of drugs, such as cardiac glycosides, antihypertensive agents and diuretics. By relieving anxiety, helps encourage productive activities. Has a wide margin of safety and, in proper maintenance dosage, seldom impairs mental acuity or ability to function. Often effective in extended therapy, usually without diminution of effect or need for increase in dosage—in protracted use, periodic blood counts and liver function tests are advisable.

References: 1. Sigler, L. H.: *Geriatrics*, 22:(9) 97, 1967.
2. Thomas, C. B.: *Johns Hopkins Med. J.*, 122:69, 1968.

Before prescribing, please consult complete product information, a summary of which follows:

Indications: Indicated when anxiety, tension and apprehension are significant components of the clinical profile.
Contraindications: Patients with known hypersensitivity to the drug.

Warnings: Caution patients about possible combined effects with alcohol and other CNS depressants. As with all CNS-

acting drugs, caution patients against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving). Though physical and psychological dependence have rarely been reported on recommended doses, use caution in administering to addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions), following discontinuation of the drug and similar to those seen with barbiturates, have been reported. Use of any drug in pregnancy, lactation, or in women of childbearing age requires that its potential benefits be weighed against its possible hazards.

Precautions: In the elderly and debilitated, and in children over six, limit to smallest effective dosage (initially 10 mg or less per day) to preclude ataxia or oversedation, increasing gradually as needed and tolerated. Not recommended in children under six. Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider individual pharmacologic effects, particularly in use of potentiating drugs such as MAO inhibitors and phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function.

Paradoxical reactions (e.g., excitement, stimulation and acute rage) have been reported in psychiatric patients and hyperactive aggressive children. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation have been reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship has not been established clinically.


Adverse Reactions: Drowsiness, ataxia and confusion may occur, especially in the elderly and debilitated. These are reversible in most instances by proper dosage adjustment, but are also occasionally observed at the lower dosage ranges. In a few instances syncope has been reported. Also encountered are isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent and generally controlled with dosage reduction; changes in EEG patterns (low-voltage fast activity) may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice and hepatic dysfunction have been reported occasionally, making periodic blood counts and liver function tests advisable during protracted therapy.

To curb anxiety in
the postcoronary patient
adjunctive

Librium®
(chlordiazepoxide HCl)

10-mg capsules

 **Roche**
LABORATORIES
Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110



**Not too little, not too much...
but just right!**

"Just right" amounts of Ilosone Liquid 250
can be dispensed easily from the pint bottle in *any* quantity
you specify to meet your patients' precise needs—
without regard to package size.

ready-mixed
Ilosone® Liquid 250

Erythromycin Estolate

(equivalent to 250 mg. of base per 5-ml. teaspoonful)

Additional information available
to the profession on request.
Eli Lilly and Company
Indianapolis, Indiana 46206



100204



Over 5 million prescriptions already filled at Macy's



*All 14 Macy Prescription Centers
satisfy your patients with
efficient professional service*



- You may phone in your patients' prescriptions (except for narcotics, amphetamines, meprobamates and barbiturates).
- Your patients may have their prescriptions delivered anywhere.

- We serve your patient with a full line of convalescent and diabetic needs.
- Your patients may charge their prescriptions on their Macy Shopping Accounts or C-T Plans.

Macy's Herald Square, OX 5-4400

Macy's Parkchester, TA 8-7000

Macy's Jamaica, OL 7-9000

Macy's Flatbush, UL 6-5000

Macy's Roosevelt Field, PI 6-8200

Macy's Huntington, AR 1-3000

Macy's White Plains, WH 6-5015

Macy's New Haven, Conn. 624-9271

Macy's Bay Shore, MO 5-8400

Macy's Queens, AR 1-9100

Macy's Colonie, 459-1950

Macy's New Rochelle, 633-7700

Macy's Smith Haven, 724-4900

Macy's Kings Plaza, 253-3100



BULLETIN of

THE NEW YORK ACADEMY OF MEDICINE

The *Bulletin* contains scientific papers presented at the Academy, including:

- ADDRESSES PRESENTED AT STATED AND ANNUAL MEETINGS
- PAPERS PRESENTED AT SECTIONS OF THE ACADEMY
- PROCEEDINGS OF SYMPOSIA AND CONFERENCES
- PUBLIC HEALTH REPORTS

Published monthly

ALL SUBSCRIPTIONS AND ORDERS FOR BACK NUMBERS SHOULD BE SENT TO:
S-H SERVICE AGENCY, INC., 866 Third Avenue, New York, N. Y. 10022

**Annual subscription: United States \$15.00, Canada \$16.00,
All other countries \$17.00 — single copies \$3.00**

Symbols in a life of
psychic tension

M.A.
in Communications

TV
director, late night
variety show

ECG
and complete
examination normal
(persistent
precordial distress)

Valium® (diazepam)

2-mg, 5-mg, 10-mg tablets
t.i.d. and *h.s.*

for relief of psychic
tension and resultant
somatic symptoms
(including tension-
induced insomnia)

... usually well tolerated

Before prescribing, please consult
complete product information, a
summary of which follows:

Indications: Tension and anxiety
states; somatic complaints which are
concomitants of emotional factors;
psychoneurotic states manifested by
tension, anxiety, apprehension,
fatigue, depressive symptoms or
agitation; acute agitation, tremor,
delirium tremens and hallucinosis
due to acute alcohol withdrawal; ad-
jectively in skeletal muscle spasm
due to reflex spasm to local pathol-
ogy, spasticity caused by upper
motor neuron disorders, athetosis,
stiff-man syndrome, convulsive
disorders (not for sole therapy).

Contraindicated: Known hypersensi-
tivity to the drug. Children under 6
months of age. Acute narrow angle
glaucoma.

Warnings: Not of value in psychotic
patients. Caution against hazardous
occupations requiring complete
mental alertness. When used ad-
jectively in convulsive disorders,

possibility of increase in frequency
and/or severity of grand mal seizures
may require increased dosage of
standard anticonvulsant medication;
abrupt withdrawal may be associated
with temporary increase in frequency
and/or severity of seizures. Advise
against simultaneous ingestion of
alcohol and other CNS depressants.
Withdrawal symptoms have occurred
following abrupt discontinuance.
Keep addiction-prone individuals
under careful surveillance because of
their predisposition to habituation
and dependence. In pregnancy, lac-
tation or women of childbearing age,
weigh potential benefit against pos-
sible hazard.

Precautions: If combined with other
psychotropics or anticonvulsants,
consider carefully pharmacology of
agents employed. Usual precautions
indicated in patients severely de-
pressed, or with latent depression,
or with suicidal tendencies. Observe
usual precautions in impaired renal
or hepatic function. Limit dosage to

smallest effective amount in elderly
and debilitated to preclude ataxia or
oversedation.

Side Effects: Drowsiness, confusion,
diplopia, hypotension, changes in
libido, nausea, fatigue, depression,
dysarthria, jaundice, skin rash,
ataxia, constipation, headache, in-
continence, changes in salivation,
slurred speech, tremor, vertigo,
urinary retention, blurred vision.
Paradoxical reactions such as acute
hyperexcited states, anxiety, halluci-
nations, increased muscle spasticity,
insomnia, rage, sleep disturbances,
stimulation, have been reported;
should these occur, discontinue
drug. Isolated reports of neutropenia,
jaundice; periodic blood counts and
liver function tests advisable during
long-term therapy.



Roche
LABORATORIES

Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110

